

IL13 Rabbit mAb [25dO]

Cat NO. :A23289

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB	Human	P35225	13kDa	Rabbit	IgG	50ul,100ul,200ul	

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

					te:		
	\sim 10		~	~	-	-	
\mathbf{u}	UH	IIU	u	а	LE	=	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human IL13

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Secreted.

Function:

Cytokine (PubMed:8096327, PubMed:8097324). Inhibits inflammatory cytokine production (PubMed:8096327). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). May be critical in regulating inflammatory and immune responses (PubMed:8096327, PubMed:8097324). Positively regulates IL31RA expression in macrophages (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

IL13 Rabbit mAb [25dO] Images



Western blot (SDS PAGE) analysis of extracts from human recombinant protein cell lysate. Using IL13 Rabbit mAb [25dO]at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com