

# SOX11 Rabbit mAb [zBK2]

Cat NO. :A25003

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB ICC/IF IP	Human,Mouse,R	P35716	70kDa	Rabbit	IgG	50ul,100ul,200ul
	at					

Applications detail:	Application	Dilution
	WB	1:1000-2000
	ICC/IF	1:100
	The optimal dilutions should be o	letermined by the end user

_				
( :n	nı	חור	gat	Δ.
$\sim$		чç	146	<b>-</b>

UnConjugate

Form:

Liquid

sensitivity:

**Endogenous** 

Purification:

Affinity-chromatography

# Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human SOX11

### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

### Tissue specificity:

Expressed primarily in the brain and heart, with low expression in the kidney, pancreas and muscle..

### Subcellular location:

Nucleus.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus Mi: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Transcription factor that acts as a transcriptional activator (PubMed:24886874). Binds cooperatively with POU3F2/BRN2 or POU3F1/OCT6 to gene promoters, which enhances transcriptional activation (By similarity). Acts as a transcriptional activator of TEAD2 by binding to its gene promoter and first intron (By similarity). Plays a redundant role with SOX4 and SOX12 in cell survival of developing tissues such as the neural tube, branchial arches and somites, thereby contributing to organogenesis (By similarity).

## **Validation Data:**

#### SOX11 Rabbit mAb [zBK2] Images



Western blot (SDS PAGE) analysis of extracts from SHSY5Y cell lysate. Using SOX11 Rabbit mAb [zBK2]at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com