

**CHX10 Rabbit mAb [EVFv]**

**Cat NO. :A37947**

**Information:**

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB FC	Human,Mouse,R at	P58304	50kDa	Rabbit	IgG	50ul,100ul,200ul

**Applications detail:**

Application	Dilution
WB	1:1000-2000
The optimal dilutions should be determined by the end user	

**Conjugate:**

UnConjugate

**Form:**

Liquid

**sensitivity:**

Endogenous

**Purification:**

Affinity-chromatography

**Specificity:**

Antibody is produced by immunizing animals with A synthesized peptide derived from human CHX10

**Storage buffer and conditions:**

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

**Tissue specificity:**

Abundantly expressed in retinal neuroblasts during eye development and in the inner nuclear layer of the adult retina. Within this layer, expression is stronger in the outer margin where bipolar

**Subcellular location:**

Nucleus.

**Function:**

**Introduction:** **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


**Cross Reactivity:** **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vr:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

**For Research Use Only. Not For Use In Diagnostic Procedures.**

Acts as a transcriptional regulator through binding to DNA at the consensus sequence 5'-[TC]TAATT[AG][AG]-3' upstream of gene promoters (PubMed:27301076). Plays a significant role in the specification and morphogenesis of the sensory retina (By similarity). Mediates differentiation of V2a interneurons by repression of motor neuron gene transcription, via competitively binding to response elements that are activated by the ISL1-LHX3 complex, such as VSX1 (PubMed:17919464, PubMed:27477290). Acts as a positive transcriptional regulator of NXNL1, regulation is significantly increased in synergy with VSX1 (By similarity). Acts as a negative transcriptional regulator of MITF (By similarity). Represses SAG transcription by competitive inhibition of ISL1-LHX3 response elements (PubMed:16236706, PubMed:27477290). Binds to the photoreceptor conserved element-1 (PCE-1) in the promoter of rod photoreceptor arrestin SAG and acts as a transcriptional repressor (By similarity). Plays a significant role in the specification and morphogenesis of the sensory retina (By similarity). Involved in the development of retinal ganglion cells (RGCs) which leads to release of SHH by RGCs, promoting Hedgehog signaling and subsequent proliferation of retinal progenitor cells (By similarity). Participates in the development of the cells of the inner nuclear layer, by promoting postnatal differentiation of bipolar cells with a comparable inhibition of rod cell differentiation (By similarity). May play a role in the maintenance of neural retina identity during development by regulation of canonical Wnt genes and CTNNB1 localization, suggesting a role in the regulation of canonical Wnt signaling (PubMed:27301076)..

## Validation Data:

### CHX10 Rabbit mAb [EVFv] Images



Western blot ( SDS PAGE ) analysis of extracts from Y79 cell lysate. Using CHX10 Rabbit mAb [EVFv] at dilution of 1:1000 incubated at 4°C over night.

View more information on <http://naturebios.com>

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.