

MTH1 Rabbit mAb [30e1]

Cat NO. :A67035

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB IHC ICC/IF IP FC	Human	P36639	18kDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application	Dilution
WB	1:1000-2000
IHC	1:100
ICC/IF	1:100
The optimal dilutions should be determined by the end user	

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Affinity-chromatography

Specificity:

Antibody is produced by immunizing animals with A synthesized peptide derived from human MTH1

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

Tissue specificity:

Widely expressed with highest expression in thymus, testis, embryo and proliferating blood lymphocytes..

Subcellular location:

[Isoform p18]: Cytoplasm, cytosol. Mitochondrion matrix. Nucleus.

Function:

Introduction: **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


Cross Reactivity: **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vr:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine
Dg: dog **Pg:** pig **Hr:** horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Oxidized purine nucleoside triphosphate hydrolase which is a prominent sanitizer of the oxidized nucleotide pool (PubMed:8226881, PubMed:7713500, PubMed:10608900, PubMed:12857738, PubMed:22556419, PubMed:26238318, PubMed:24695224, PubMed:24695225, PubMed:28679043). Catalyzes the hydrolysis of 2-oxo-dATP (2-hydroxy-dATP) into 2-oxo-dAMP (PubMed:10373420). Has also a significant hydrolase activity toward 2-oxo-ATP, 8-oxo-dGTP and 8-oxo-dATP (PubMed:10373420, PubMed:11139615). Through the hydrolysis of oxidized purine nucleoside triphosphates, prevents their incorporation into DNA and the subsequent transversions A:T to C:G and G:C to T:A (PubMed:8226881, PubMed:10373420, PubMed:10608900, PubMed:11756418, PubMed:12857738, PubMed:16607562, PubMed:24695224, PubMed:24695225, PubMed:26999531, PubMed:28035004). Also catalyzes the hydrolysis of methylated purine nucleoside triphosphate preventing their integration into DNA (PubMed:30304478, PubMed:32144205). Through this antimutagenic activity protects cells from oxidative stress (PubMed:8226881, PubMed:7713500, PubMed:10608900, PubMed:12857738, PubMed:24695224, PubMed:24695225, PubMed:30304478, PubMed:32144205)..

Validation Data:

MTH1 Rabbit mAb [3Oe1] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cell lysate.Using MTH1 Rabbit mAb [3Oe1]at dilution of 1:1000 incubated at 4°C over night.

View more information on <http://naturebios.com>

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.