

ENTPD5 Rabbit mAb [0KEF]

Cat NO. :A18007

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	O75356	48KDa	Rabbit	IgG	50ul,100ul,200ul

Applications detail:

Application

WB

1:1000-2000

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human ENTPD5.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Expressed in adult liver, kidney, prostate, testis and colon. Much weaker expression in other tissues..

Subcellular location:

Endoplasmic reticulum. Secreted.

Function:

Hydrolyzes nucleoside diphosphates with a preference for GDP, IDP and UDP compared to ADP and CDP (PubMed:15698960, PubMed:10400613). In the lumen of the endoplasmic reticulum, hydrolyzes UDP that acts as an end-product feedback inhibitor of the UDP-Glc:glycoprotein glucosyltransferases. UMP can be transported back by an UDP-sugar antiporter to the cytosol where it is consumed to regenerate UDP-glucose. Therefore, it positively regulates protein reglucosylation by clearing UDP from the ER lumen and by promoting the regeneration of UDP-glucose. Protein reglucosylation is essential to proper glycoprotein folding and quality control in the ER (By similarity)...

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

ENTPD5 Rabbit mAb [0KEF] Images



Western blot (SDS PAGE) analysis of extracts from Mouse liver Tissue lysate. Using ENTPD5 Rabbit mAb [0KEF] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com