TROP2 Rabbit mAb [8675]

Cat NO. :A21819

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M	P09758	37-50 kDa	Rabbit	lgG	100ul,200ul

Applications detail:

Application	Dilution		
WB	1:1000-2000		
ІНС	1:100		
ICC/IF	1:100		
The optimal dilutions should be determined by the end user			

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human TROP2

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol $\ (buffer) \ .$

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Placenta, pancreatic carcinoma cell lines.

Subcellular location:

Membrane, Single-pass type I membrane protein.

Function:

May function as a growth factor receptor.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Validation Data:

TROP2 Rabbit mAb [8675] Images



Western blot (SDS PAGE) analysis of extracts from HCT 116 cells.Using TROP2Rabbit mAb [8675] at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.