

ABAT/GABA-T Rabbit mAb [4KP9]

Cat NO. :A44833

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	H,M,R	P80404	50 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
IHC	1:100			
ICC/IF	1:100			
The optimal dilutions should be determined by the end user				

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human ABAT/GABA-T

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Liver > pancreas > brain > kidney > heart > placenta..

Subcellular location:

Mitochondrion matrix.

Function:

Catalyzes the conversion of gamma-aminobutyrate and L-beta-aminoisobutyrate to succinate semialdehyde and methylmalonate semialdehyde, respectively (PubMed:10407778, PubMed:15528998). Can also convert delta-aminovalerate and beta-alanine (By similarity)..

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

ABAT/GABA-T Rabbit mAb [4KP9] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells .Using ABAT/GABA-TRabbit mAb [4KP9] at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night.

View more information on http://naturebios.com