## NEUROD1 Mouse mAb[4WKP]

Cat NO. :A45422

## Information:

Ар	plications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
	WB,IHC	H,M	Q13562	40kda	Mouse	lgG	50ul 100ul,200ul

### **Applications detail:**

Application	Dilution			
WB	1:1000-2000			
IHC	1:100			
The optimal dilutions should be determined by the end user				

### Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

## **Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of human NEUROD1.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

**Tissue specificity:** 

## Subcellular location:

Cytoplasm. Nucleus.

#### **Function**:

Acts as a transcriptional activator: mediates transcriptional activation by binding to E box-containing promoter consensus core sequences 5'-CANNTG-3'. Associates with the p300/CBP transcription coactivator complex to stimulate transcription of the secretin gene as well as the gene encoding the cyclin-dependent kinase inhibitor CDKN1A. Contributes to the regulation of several cell differentiation pathways, like those that promote the formation of early retinal ganglion cells, inner ear sensory neurons, granule cells forming either the cerebellum or the dentate gyrus cell layer of the hippocampus, endocrine islet cells of the pancreas and enteroendocrine cells of the small intestine. Together with PAX6 or SIX3, is required for the regulation of amacrine cell fate

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

specification. Also required for dendrite morphogenesis and maintenance in the cerebellar cortex. Associates with chromatin to enhancer regulatory elements in genes encoding key transcriptional regulators of neurogenesis (By similarity)..

# Validation Data:

## NEUROD1 Mouse mAb[4WKP] Images



Western blot (SDS PAGE) analysis of extracts from Y79 cells.Using NEUROD1 Mouse mAb IgG [4WKP] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.