Junctional Adhesion Molecule 1/JAM-A Rabbit mAb [I80L]

Cat NO. :A50534

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	н	Q9Y624	33 kDa	Rabbit	lgG	100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
ІНС	1:100			
The optimal dilutions should be	e optimal dilutions should be determined by the end user			

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human Junctional Adhesion Molecule 1/JAM-A

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed in endothelium, epithelium and leukocytes (at protein level)..

Subcellular location:

Cell junction, tight junction. Cell membrane,Single-pass type I membrane protein.

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

Seems to play a role in epithelial tight junction formation. Appears early in primordial forms of cell junctions and recruits PARD3 (PubMed:11489913). The association of the PARD6-PARD3 complex may prevent the interaction of PARD3 with JAM1, thereby preventing tight junction assembly (By similarity). Plays a role in regulating monocyte transmigration involved in integrity of epithelial barrier (By similarity). Ligand for integrin alpha-L/beta-2 involved in memory T-cell and neutrophil transmigration (PubMed:11812992). Involved in platelet activation (PubMed:10753840)..., (Microbial infection) Acts as a receptor for Mammalian reovirus sigma-1..., (Microbial infection) Acts as a receptor for Human Rotavirus strain Wa..

Validation Data:

Junctional Adhesion Molecule 1/JAM-A Rabbit mAb [I80L] Images



Western blot(SDS PAGE) analysis of extracts from HeLa cells cell lysate.Using Junctional Adhesion Molecule 1/JAM-ARabbit mAb [I80L] at dilution of 1:1000 incubated

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.