Vasorin Rabbit mAb [4YF8]

Cat NO. :A53741

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	Q6EMK4	72 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human Vasorin.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Expressed at highest levels in aorta, at intermediate levels in kidney and placenta and at lowest levels in brain,

heart, liver, lung and skeletal muscle. Within the aorta, the strongest expression

Subcellular location:

Membrane, Single-pass type I membrane protein. Secreted.

Function:

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry


Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

May act as an inhibitor of TGF-beta signaling..

Validation Data:

Vasorin Rabbit mAb [4YF8] Images



Western blot (SDS PAGE) analysis of extracts from Human fetal heart tissue lyaste.using Vasorin Rabbit mAb [4YF8] at dilution of 1:1000 incubated at 4° C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.