

**MYSM1 Rabbit mAb [7621]**

**Cat NO. :A54153**

**Information:**

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	H,M,R	Q5VVJ2	66kd kDa	Rabbit	IgG	100ul,200ul

**Applications detail:**

Application	Dilution
WB	1:1000-2000
The optimal dilutions should be determined by the end user	

**Conjugate:**

UnConjugate

**Form:**

Liquid

**sensitivity:**

Endogenous

**Purification:**

Protein A purification

**Specificity:**

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human MYSM1

**Storage buffer and conditions:**

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at -20°C or -80°C.

Products are valid for one natural year of receipt.Avoid repeated freeze / thaw cycles.

**Tissue specificity:**

**Subcellular location:**

Nucleus. Cytoplasm.

**Function:**

Metalloprotease with deubiquitinase activity that plays important regulator roles in hematopoietic stem cell function, blood cell production and immune response (PubMed:24062447, PubMed:26220525, PubMed:28115216). Participates in the normal programming of B-cell responses to antigen after the maturation process (By similarity). Within the cytoplasm, plays critical roles in the repression of innate immunity and autoimmunity (PubMed:33086059). Removes 'Lys-63'-linked polyubiquitins from TRAF3 and TRAF6 complexes (By similarity). Attenuates NOD2-mediated inflammation and tissue injury by promoting 'Lys-63'-linked deubiquitination of RIPK2 component (By similarity). Suppresses the CGAS-STING1 signaling pathway by

**Introduction:** **WB:** Western Blot **IP:** Immunoprecipitation **IHC:** Immunohistochemistry **ChIP:** Chromatin Immunoprecipitation **ICC/IF:** Immunocytochemistry/Immunofluorescence **F:** Flow Cytometry


**Cross Reactivity:** **H:** human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Ml:** mink **C:** chicken **Dm** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Hr:** horse

**For Research Use Only. Not For Use In Diagnostic Procedures.**

cleaving STING1 'Lys-63'-linked ubiquitin chains (PubMed:33086059). In the nucleus, acts as a hematopoietic transcription regulator derepressing a range of genes essential for normal stem cell differentiation including EBF1 and PAX5 in B-cells, ID2 in NK-cell progenitor or FLT3 in dendritic cell precursors (PubMed:24062447). Deubiquitinates monoubiquitinated histone H2A, a specific tag for epigenetic transcriptional repression, leading to dissociation of histone H1 from the nucleosome (PubMed:17707232)..

## Validation Data:

### MYSM1 Rabbit mAb [7621] Images



Western blot (SDS PAGE) analysis of extracts from HEK-293 cells. Using MYSM1 Rabbit mAb [7621] at dilution of 1:1000 incubated at 4°C over night.

View more information on <http://naturebios.com>

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.