

ATOX1 Rabbit mAb [A9X2]

Cat NO. :A55980

Information:

| Applications | Reactivity: | UniProt ID: | MW(kDa) | Host | Isotype | Size | |
|--------------|-------------|-------------|---------|--------|---------|-------------|--|
| WB,IHC | н | O00244 | 7 kDa | Rabbit | IgG | 100ul,200ul | |

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

| | | | | | te: | | |
|--------------|-----------|-----|---|---|-----|---|--|
| | \sim 10 | | ~ | ~ | - | - | |
| \mathbf{u} | UH | IIU | u | а | LE | = | |
| | | | | | | | |

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human ATOX1

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Ubiquitous.

Subcellular location:

Function:

Binds and deliver cytosolic copper to the copper ATPase proteins. May be important in cellular antioxidant defense.


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

ATOX1 Rabbit mAb [A9X2] Images



Western blot (SDS PAGE) analysis of extracts from HepG2 cells.Using ATOX1Rabbit mAb [A9X2] at dilution of 1:1000 incubated at 4° C over night.

View more information on http://naturebios.com