

SSB Rabbit mAb [7345]

Cat NO. :A60044

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size	
WB,IHC	H,M,R	P05455	47 kDa	Rabbit	IgG	50ul,100ul,200ul	

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

					te:		
	\sim 10		~	~	-	-	
\mathbf{u}	UH	IIU	u	а	LE	=	

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human SSB.

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Subcellular location:

Nucleus.

Function:

Binds to the 3' poly(U) terminus of nascent RNA polymerase III transcripts, protecting them from exonuclease digestion and facilitating their folding and maturation (PubMed:3192525, PubMed:2470590). In case of Coxsackievirus B3 infection, binds to the viral internal ribosome entry site (IRES) and stimulates the IRES-mediated translation (PubMed:12384597)..

Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

SSB Rabbit mAb [7345] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cells lyastes.using SSB Rabbit mAb [7345] at dilution of 1:1000 incubated at $4^{\circ}\mathrm{C}$ over night

View more information on http://naturebios.com