# BLBP Rabbit mAb [Q4FV]

Cat NO. :A65549

## Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	O15540	14 kDa	Rabbit	lgG	100ul,200ul

## **Applications detail:**

Application	Dilution		
WB	1:1000-2000		
ІНС	1:100		
The optimal dilutions should be	he optimal dilutions should be determined by the end user		

## Conjugate:

UnConjugate

Form:

Liquid

#### sensitivity:

Endogenous

## **Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human BLBP

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

## **Tissue specificity:**

Expressed in brain and other neural tissues.

# Subcellular location:

Cytoplasm.

#### **Function**:

B-FABP could be involved in the transport of a so far unknown hydrophobic ligand with potential morphogenic activity during CNS development. It is required for the establishment of the radial glial fiber system in developing brain, a system that is necessary for the migration of immature neurons to establish cortical layers (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cvtometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

# Validation Data:

## BLBP Rabbit mAb [Q4FV] Images



Western blot (SDS PAGE) analysis of extracts from Mouse brain.Using BLBPRabbit mAb [Q4FV] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.