

UFM1 Rabbit mAb [2M48]

Cat NO. :A65602

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	P61960	9 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human UFM1

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$

Tissue specificity:

Subcellular location:

Nucleus. Cytoplasm.

Function:

Ubiquitin-like modifier which can be covalently attached via an isopeptide bond to lysine residues of substrate proteins as a monomer or a lysine-linked polymer (PubMed:15071506, PubMed:20018847, PubMed:29868776, PubMed:27653677). The so-called ufmylation, requires the UFM1-activating E1 enzyme UBA5, the UFM1-conjugating E2 enzyme UFC1, and the UFM1-ligase E3 enzyme UFL1 (PubMed:15071506, PubMed:20018847, PubMed:29868776, PubMed:27653677). Ufmylation is involved in reticulophagy (also called ER-phagy) induced in response to endoplasmic reticulum stress (PubMed:32160526). Ufmylation of TRIP4 regulates nuclear receptors-mediated transcription (PubMed:25219498)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

UFM1 Rabbit mAb [2M48] Images



Western blot (SDS PAGE) analysis of extracts from SH-SY5Y cells.Using UFM1Rabbit mAb [2M48] at dilution of 1:1000 incubated at 4° over night.

View more information on http://naturebios.com