

NCE2/UBE2F Rabbit mAb [D6X1]

Cat NO. :A67212

Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC,ICC/IF	н	Q969M7	24 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application	Dilution			
WB	1:1000-2000			
IHC	1:100			
ICC/IF	1:100			
The optimal dilutions should be determined by the end user				

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human NCE2/UBE2F

Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

Tissue specificity:

Widely expressed (at protein level)..

Subcellular location:

Function:

Accepts the ubiquitin-like protein NEDD8 from the UBA3-NAE1 E1 complex and catalyzes its covalent attachment to other proteins. The specific interaction with the E3 ubiquitin ligase RBX2, but not RBX1, suggests that the RBX2-UBE2F complex neddylates specific target proteins, such as CUL5..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



Validation Data:

NCE2/UBE2F Rabbit mAb [D6X1] Images



Western blot (SDS PAGE) analysis of extracts from Jurkat cells.Using NCE2/UBE2FRabbit mAb [D6X1] at dilution of 1:1000 incubated at $4\,^\circ\!\mathrm{C}$ over night.

View more information on http://naturebios.com