

# Sp7 / Osterix Rabbit mAb [8E66]

Cat NO. :A83446

#### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB,IHC	H,M,R	Q8TDD2	45, 47 kDa	Rabbit	IgG	100ul,200ul

Applications detail:

Application

WB

1:1000-2000

IHC

The optimal dilutions should be determined by the end user

Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

Purification:

Protein A purification

Specificity:

Antibody is produced by immunizing animals with a synthetic peptide at the sequence of human Sp7 / Osterix

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol (buffer) .

Shipped at 4°C. Store at-20°C or -80°C.

 $\label{products} \textbf{Products are valid for one natural year of receipt.} \textbf{Avoid repeated freeze} \ \textit{I} \ \textbf{thaw cycles}.$ 

### Tissue specificity:

Restricted to bone-derived cell..

### Subcellular location:

Nucleus.

#### Function:

Transcriptional activator essential for osteoblast differentiation (PubMed:23457570). Binds to SP1 and EKLF consensus sequences and to other G/C-rich sequences (By similarity)..


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/
Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse



## **Validation Data:**

## Sp7 / Osterix Rabbit mAb [8E66] Images



Western blot(SDS PAGE) analysis of extracts from Saos-2 cells .Using Sp7 / OsterixRabbit mAb [8E66] at dilution of 1:1000 incubated at  $4^{\circ}$ C over night.

View more information on http://naturebios.com