# C16orf88 Rabbit mAb [44l6]

Cat NO. :A95739

### Information:

Applications	Reactivity:	UniProt ID:	MW(kDa)	Host	Isotype	Size
WB	н	Q1ED39	52 kda	Rabbit	lgG	100ul,200ul

#### **Applications detail:**

# Application Dilution WB 1:1000-2000 The optimal dilutions should be determined by the end user

#### Conjugate:

UnConjugate

Form:

Liquid

sensitivity:

Endogenous

## **Purification**:

Protein A purification

#### Specificity:

Antibody is produced by immunizing animals with a synthetic peptide of Human C16orf88.

#### Storage buffer and conditions:

Antibody store in 10 mM PBS, 0.5mg/ml BSA, 50% glycerol  $\ (buffer) \ .$ 

Shipped at 4°C. Store at-20°C or -80°C.

Products are valid for one natural year of receipt. Avoid repeated freeze / thaw cycles.

**Tissue specificity:** 

#### Subcellular location:

Nucleus, nucleolus.

**Function**:


Introduction: WB: Western Blot IP: Immunoprecipitation IHC: Immunohistochemistry ChIP: Chromatin Immunoprecipitation ICC/IF: Immunocytochemistry/ Immunofluorescence F: Flow Cytometry

Cross Reactivity: H: human M: mouse R: rat Hm: hamster Mk: monkey Vir: virus MI: mink C: chicken Dm D. melanogaster X: Xenopus Z: zebrafish B: bovine Dg: dog Pg: pig Hr: horse

For Research Use Only. Not For Use In Diagnostic Procedures.

# Validation Data:

## C16orf88 Rabbit mAb [4416] Images



Western blot (SDS PAGE) analysis of extracts from Human testis tissue lyaste.using C16orf88 Rabbit mAb [4416] at dilution of 1:1000 incubated at 4°C over night

View more information on http://naturebios.com

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 1% w/v Milk, 1X TBST at 4°C overnight.